

Accretion into the Central Cavity of a Circumbinary Disk

Daniel J. D’Orazio¹, Zoltán Haiman¹, Andrew MacFadyen^{2*}

¹*Department of Astronomy, Columbia University, 550 West 120th Street, New York, NY 10027*

²*Center for Cosmology and Particle Physics, Physics Department, New York University, New York, NY 10003*

3 October 2012

ABSTRACT

A near-equal mass binary black hole can clear a central cavity in a circumbinary accretion disk; however, previous works have revealed accretion streams entering this cavity. Here we use 2D hydrodynamical simulations to study the accretion streams and their periodic behavior. In particular, we perform a suite of simulations, covering different binary mass ratios $q = M_2/M_1$ in the range $0.01 \leq q \leq 1$. In each case, we follow the system for several thousand binary orbits, until it relaxes to a stable accretion pattern. We find the following results: (i) while the binary is efficient in maintaining a low-density cavity, the time-averaged mass accretion rate into the cavity, through narrow coherent accretion streams, is suppressed by at most a factor of ~ 5 , compared to a disk with a single BH with the same mass; (ii) the largest suppression occurs for $q \approx 0.05$; binaries whose mass ratios are either lower or higher both suppress accretion less significantly; (iii) for $q \gtrsim 0.05$, the accretion rate is strongly modulated by the binary, and depending on the precise value of q , the power spectrum of the accretion rate shows either one, two, or three distinct periods; and (v) for $q \lesssim 0.05$, the accretion rate becomes steady, with no time-variations. Most binaries produced in galactic mergers are expected to have $q \gtrsim 0.05$. If the luminosity of these binaries tracks their accretion rate, then a periodogram of their light-curve could help in their identification, and to constrain their mass ratio and disk properties.

Key words: black hole physics — accretion, accretion discs — galaxies: active — gravitational waves

1 INTRODUCTION

Massive black holes (MBHs) appear to reside in the nuclei of most nearby galaxies (see, e.g., reviews by Kormendy & Richstone 1995 and Ferrarese & Ford 2005). In hierarchical structure formation models, galaxies are built up by mergers between lower-mass progenitors, which deliver nuclear MBHs (e.g. Springel et al. 2005; Robertson et al. 2006), along with a significant amount of gas (Barnes & Hernquist 1992), to the central region of the newly born post-merger galaxy. Since mergers are common (e.g. Haehnelt & Kauffmann 2002), it follows that massive black hole binaries (MBHBs) should also be common in galactic nuclei.

Despite this expectation, observational evidence for MBHBs remains scarce (see, e.g., a review by Komossa 2006). The dearth of MBHBs could be attributed to several factors: it is possible that typically only one of the two BHs is active at spatially resolvable separations; binaries may also lose their angular momentum efficiently due to the surrounding stars and gas and quickly move to spatially un-

resolvable orbital separations. Another possible hindrance, which we address in this paper, is that the outward gravitational torques from the binary can balance the inward viscous and pressure torques, clearing a central cavity in a putative circumbinary gas disk (Artymowicz & Lubow 1994), possibly rendering the system too dim for detection. Overall, identifying MBHBs is difficult, and a better understanding of their expected observational signatures, especially those based on time-variability (Haiman et al. 2009), is needed. Merging MBHBs should be unambiguously identifiable by gravitational wave (GW) detectors, such as eLISA (Amaro-Seoane & et. al 2012) or ongoing Pulsar Timing Arrays (e.g. Lommen 2012). Identifying the electromagnetic (EM) counterparts of these GW sources (among the many false candidate galaxies in the GW error box) will, however, likewise require an understanding of their observational signatures.

Recent studies have explored the gasdynamics of circumbinary accretion disks around near-equal mass binaries in some detail. Since the system is not axisymmetric, this requires a two- or three-dimensional treatment. MacFadyen & Milosavljević (2008) (hereafter MM08) have run two-dimensional hydrodynamical simulations for an equal mass binary. Three-dimensional smoothed particle hydrodynam-

* dorazio@astro.columbia.edu; zoltan@astro.columbia.edu; macfadyen@nyu.edu

ical (SPH) simulations have been carried out for equal-mass and 2:1 mass-ratio binaries by Hayasaki et al. (2007) and for a 3:1 mass-ratio binary by Cuadra et al. (2009) and Roedig et al. (2012). Shi et al. (2011) have followed up on the work of MM08 for an equal-mass binary by running 3D magneto-hydrodynamical (MHD) simulations, and Noble et al. (2012) have further added a post-Newtonian treatment of general relativistic (GR) effects, and followed the disk through the late stages of orbital inspiral (from orbital separation $r = 20M$ to $8M$). Farris et al. (2011) followed the merger of an equal-mass binary and a surrounding disk through merger in full 3D general relativistic MHD, starting from $10M$. Finally, Farris et al. (2012) have added gas-cooling to GRMHD simulations of an equal-mass binary prior to decoupling, through decoupling and to merger starting from $10M$. A generic result of all of these studies is that a low-density cavity is carved out by the binary torque, but gas leaks into the cavity through non-axisymmetric streams. These streams can power significant accretion onto the binary components, and should lead to bright EM emission.

A particularly promising feature is that the accretion rate onto the BHs can be both high and strongly variable, modulated by the binary’s orbital motion. This could allow a detection of sub-pc binaries by looking for periodic variations in the luminosity of AGN-like objects (Haiman et al. 2009) or periodic shifts and intensity variations of spectral lines (e.g. Haiman et al. 2009; Shen & Loeb 2010; Eracleous et al. 2011 and references therein). If the accretion remains significant and periodic down to $\ll$ pc separations, then it could also enable the identification of EM counterparts of gravitational wave sources: either for precursors to eLISA sources in the $M = 10^5 - 10^7 M_\odot$ range (Kocsis et al. 2006, 2008) or by detecting periodic modulations of more massive $M = 10^8 - 10^9 M_\odot$ binaries discovered by pulsar timing arrays (PTAs; Tanaka et al. 2012; Sesana et al. 2012).

Although existing studies have focused on near-equal mass MBHBs, in reality, coalescing MBHBs should have a distribution of mass ratios $q \equiv M_2/M_1$.¹ Mergers occur between galaxies over a wide range of sizes, harboring central BHs of different masses, so that MBHBs resulting from galactic mergers should have a correspondingly wide range of mass ratios. Studies based on Monte-Carlo realizations of dark matter merger trees indeed find broad distributions between $10^{-2} \lesssim q < 1$, generally peaking in the range $q \sim 0.1 - 1$ (e.g. Volonteri et al. 2003; Sesana et al. 2005, 2012; Gergely & Biermann 2012). However, the predictions depend on the occupation fraction of MBHs, the redshift-evolution of the correlation between the masses of MBHs and their host galaxies, as well as on the limit on the mass ratio of host galaxies whose nuclear MBHs can coalesce; $q < 0.1$ mergers could in fact be most common (e.g. Lippai et al. 2009).

Here we follow up on the earlier work of MM08 and move beyond the near-equal mass binary case. We study the periodicity and the time-averaged rate of accretion across the central cavity, by running 2D hydrodynamical simulations of a circumbinary disk for 12 different binary mass ratios ranging from $q = 0.01$ to $q = 1$. Clearly, one expects

that in the limit $q \rightarrow 0$, the accretion rate approaches that of an accretion disk around a single BH, and will no longer be time-variable. The main goal in this paper is to answer the following basic questions: *How does the mean accretion rate, and its fluctuations, depend on the mass ratio? In particular, down to what mass ratio is the mean accretion and/or its variability significantly affected by the binary torques?*

The rest of this paper is organized as follows: In § 2, we describe the setup of our numerical simulations, including a few changes we made to the public version of the Eulerian grid code FLASH, basic tests of the modified code, and the initial and boundary conditions we adopted. In § 3, we discuss scaling the simulations to physical parameters, such as black hole mass and orbital separation, and discuss the corresponding orbital and residence times. In § 4, we present our main results, where we find four distinct patterns for the time-variability of the accretion rate, as a function of the mass ratio q . We also discuss the interpretation of these results, as well as some caveats, in this section. Finally, in § 5 we conclude by briefly summarizing our main results and their implications.

2 DETAILS OF NUMERICAL SIMULATIONS

To simulate a gas disk in the gravitational field of a binary, we use the Eulerian grid-based hydrodynamical code FLASH (Version 3.2; Fryxell et al. 2000).² FLASH solves the volume-integrated fluid equations by solving the Riemann ‘shock-tube’ problem at each cell boundary. A piece-wise parabolic representation of the fluid variables is used to interpolate between cells, i.e. FLASH is a PPM code, accurate to 2nd order in both space and time. FLASH uses a monotonicity constraint, rather than artificial viscosity, to control oscillations near discontinuities. This makes it well suited for following supersonic fluid dynamics in the inner regions of circumbinary disk. FLASH also supports polar coordinates, which is convenient for simulating disks.

2.1 Numerical Implementation and Assumptions

We assume a geometrically thin accretion flow onto the binary. This permits a decoupling of the fluid equations in the z direction, perpendicular to the plane of the disk, so that we can define height-integrated fluid variables and set up simulations in two dimensions. In follow-up studies, we plan to extend these simulations to the full three dimensions, which we expect will be important in determining the amount of inflow into a putative circum-binary cavity (for recent 3D grid-based simulations, see Shi et al. 2011 and Noble et al. 2012). In the present study, we choose 2D polar coordinates (r, ϕ) and employ FLASH to solve the following standard

¹ Nixon et al. (2011) explored a range of mass ratios, using 3D SPH simulations, but restricted their study to *retrograde* disks.

² We note that MM08 used an earlier release, Version 2, of the same code.

set of 2D hydrodynamical equations:

$$\begin{aligned} \frac{\partial \Sigma}{\partial t} + \nabla \cdot (\Sigma \vec{v}) &= 0 \\ \frac{\partial \vec{v}}{\partial t} + (\vec{v} \cdot \nabla) \vec{v} &= -\frac{1}{\Sigma} \nabla P - \nabla \Phi_{\text{bin}} + \nabla \cdot \nu \nabla \vec{v} \\ \frac{\partial (\Sigma E)}{\partial t} + \nabla \cdot [(\Sigma E + P) \vec{v}] &= \Sigma \vec{v} \cdot (-\nabla \Phi_{\text{bin}}) \end{aligned} \quad (1)$$

Here Σ is the vertically integrated disk surface density, $\vec{v} = v_r \hat{r} + v_\phi \hat{\phi}$ is the fluid velocity, P is the pressure, ν is the coefficient of kinematic viscosity, E is the total internal plus kinetic energy of the fluid, $E = \epsilon + \frac{1}{2} |\vec{v}|^2$, and Φ_{bin} is the gravitational potential of the binary. The gravitational potential is inserted into the simulation by hand, and is given by

$$\begin{aligned} \Phi_{\text{bin}}(r, \phi) &= -\frac{GM(1+q)^{-1}}{\left[r^2 + \left(\frac{a}{1+q^{-1}} \right)^2 - \frac{2ra}{1+q^{-1}} \cos(\phi - \Omega_{\text{bin}} t) \right]^{1/2}} \\ &\quad - \frac{GM(1+q^{-1})^{-1}}{\left[r^2 + \left(\frac{a}{1+q} \right)^2 + \frac{2ra}{1+q} \cos(\phi - \Omega_{\text{bin}} t) \right]^{1/2}}. \end{aligned} \quad (2)$$

Here $\Omega_{\text{bin}} = (GM/a^3)^{1/2}$ is the binary's orbital frequency, a is the separation of the binary, $M_p > M_s$ are the masses of the primary and the secondary, $M = M_p + M_s$ is the total mass, and $q = M_s/M_p \leq 1$ is the mass ratio. The origin of the coordinate system is chosen coincide with the binary's center of mass. Note that we do not evolve the orbital parameters of the binary (this assumption is justified for our physical parameter choices; see discussion in § 3 below).

We neglect self-gravity of the disk. Given the local sound speed c_s , the Toomre parameter $Q \equiv c_s \Omega / (\pi G \Sigma)$ can be written as $Q \sim (H/r)(M/M_d)$, for a disk with mass M_d and vertical scale-height H in hydrostatic equilibrium. For a thin disk $H/r \lesssim 0.1$, but in all of our simulations, we choose $M \gg M_d$ and thus $Q \gg 1$, making the disk stable to gravitational fragmentation. This is justified for standard Shakura-Sunyaev disks, when the simulations are scaled to physical BH masses and sufficiently small separations (see § 3 below). For comparison, we note that in their SPH simulations, Cuadra et al. (2009) and Roedig et al. (2012) studied more massive disks, with $M_d \sim 0.4M$, making self-gravity important.

The pressure is given, as in MM08, by a locally isothermal equation of state,

$$P = c_s^2(r) \Sigma \quad (3)$$

where the sound speed is assumed to scale with radius as $c_s \propto r^{-1/2}$. For a Keplerian potential, this corresponds to a constant disk scale-height to radius ratio, $H/r = c_s/v_\phi \equiv \mathcal{M}^{-1}$, where v_ϕ is the orbital velocity in the disk and $\mathcal{M}$ is the corresponding Mach number. Throughout our simulations, we chose the disk sound speed such that, for a Keplerian azimuthal velocity, $H/r = 0.1$ (or $\mathcal{M} = 10$) everywhere.³

³ Since we actually add the binary's quadrupole potential, and the non-zero pressure makes the azimuthal velocities slightly sub-Keplerian, in practice $\mathcal{M}$ in the simulations approaches ≈ 18 near $r = a$ and becomes a constant $\mathcal{M} \approx 10$ at $r > 5a$.

To incorporate viscosity, FLASH calculates the momentum flux across cell boundaries due to viscous dissipation (the last term in the second of equations 1). To compute ν we adopt the α prescription, $\nu = \alpha c_s H$ (Shakura & Sunyaev 1973), where α is a dimensionless parameter indicating the scale of turbulent cells, and the scale height is computed from $H = c_s r / v_\phi = 0.1r$ with v_ϕ being the Keplerian value. Following MM08, we chose a constant $\alpha = 0.01$ (but see discussion in § 2.4 below). The viscosity implementation in FLASH3 follows the method of Edgar (2006), and is very different from that of FLASH2. Since neither implementation is fully supported in polar coordinates, we made adjustments to the routines that compute the momentum flux and viscous diffusion time from ν , originally in Cartesian coordinates. These changes amount to keeping track of volume elements when taking azimuthal derivatives. The method is outlined in polar coordinates in Edgar (2006). We test our viscosity implementation in § 2.4 below.

2.2 Numerical Parameter Choices

Unless specified otherwise, the inner edge of the computational domain is chosen at $r_{\text{min}} = a$ and the outer edge at $r_{\text{max}} = 100a$. Although we are only interested in the inner few r/a of the disk, extending the computations to larger radii acts as a buffer for small initial numerical transients, and also provides the reservoir of gas from which the inner regions can be fed. As in other Eulerian codes, FLASH uses a boundary zone of 'guard cells' which enforces boundary conditions. We chose a diode-type inner boundary condition: the values of fluid variables in cells bordering the guard cells are copied into the guard cells, with the restriction that no fluid enter the domain, $v_r(r_{\text{min}}) \leq 0$. We adopt an 'outflow', outer boundary condition: this is identical to the diode version, except that flow is allowed both into and out of the domain. In practice, with our initial conditions in § 2.3, we find an outflow at the outer boundary of our disk. However, the simulation does not run for a significant fraction of a viscous time at the outer radius (see below) and we expect inflow at the outer boundary to establish itself eventually.

The spatial resolution was fixed throughout the grid, with the default values set at $[\Delta r/a, \Delta \phi/2\pi] \simeq [0.024, 0.0078]$, corresponding to a grid of $\approx 4096 \times 128$ cells. The wavelength of a density wave due to a Lindblad resonance is of order $2\pi H \sim 0.6r$ (e.g. Dong et al. 2011; Duffell & MacFadyen 2012); as in MM08, the radial resolution was chosen to resolve this length scale with many cells. To test numerical convergence (see § 4.3.3 below), we performed several additional runs, increasing the radial and azimuthal resolutions, by factors of 1.42 and $1.42^2 \approx 2.0$, from the lowest resolution runs. The time resolution is set to be half of the shortest propagation time (viscous or dynamical) across a cell. The effects of changing the resolution will be discussed in § 4.3.3 below.

We run our simulations for between 4×10^3 and 10^4 binary orbits. For reference, we note that the viscous time can be related to the orbital time as

$$t_{\text{visc}} = \frac{2}{3} \frac{r^2}{\nu} = \frac{\mathcal{M}^2}{3\pi\alpha} t_{\text{orb}} \simeq 1060 \left(\frac{\mathcal{M}^2}{100} \right) \left(\frac{0.01}{\alpha} \right) t_{\text{orb}}. \quad (4)$$

Thus, our typical run of 4000 orbits corresponds to ~ 4

Table 1. Summary of our simulation runs. Low, medium, and high radial and azimuthal resolutions are denoted by “Lo Δr ”; “Mid Δr ”; “Hi Δr ” and “Lo $\Delta\phi$ ”; “Mid $\Delta\phi$ ”; “Hi $\Delta\phi$ ” and correspond to $\Delta r/a = 0.035, 0.24, 0.017$ and $\Delta\phi/2\pi = 0.0078, 0.0052, 0.0039$, respectively. The viscosity parameter α is set to 0.01 unless otherwise specified.

Mass ratio q	Spatial Resolution $[\Delta r, \Delta\phi]$	No. Orbits ($N_{\text{orb}} = t_{\text{max}}\Omega_{\text{bin}}/2\pi$)
1.0	[Lo Δr , Lo $\Delta\phi$], [Mid Δr , Lo $\Delta\phi$]	5000, 5000
1.0 ($\alpha = 0.05$)	[Lo Δr , Lo $\Delta\phi$], [Mid Δr , Lo $\Delta\phi$]	4500, 4500
0.75	[Lo Δr , Lo $\Delta\phi$], [Mid Δr , Lo $\Delta\phi$]	4500, 4500
0.5	[Lo Δr , Lo $\Delta\phi$], [Mid Δr , Lo $\Delta\phi$]	4500, 4500
0.25	[Lo Δr , Lo $\Delta\phi$], [Mid Δr , Lo $\Delta\phi$]	4800
0.1	[Lo Δr , Lo $\Delta\phi$], [Mid Δr , Lo $\Delta\phi$], [Mid Δr , Mid $\Delta\phi$], [Hi Δr , Hi $\Delta\phi$],	4500, 4500, 4500, 4200
0.075	[Lo Δr , Lo $\Delta\phi$], [Mid Δr , Lo $\Delta\phi$]	4500, 4500
0.065	[Lo Δr , Lo $\Delta\phi$]	4500
0.06	[Lo Δr , Lo $\Delta\phi$]	8000
0.05	[Lo Δr , Lo $\Delta\phi$], [Mid Δr , Lo $\Delta\phi$]	4500, 5000
0.04	[Mid Δr , Lo $\Delta\phi$]	4500
0.03	[Mid Δr , Lo $\Delta\phi$]	4500
0.01	[Lo Δr , Lo $\Delta\phi$], [Mid Δr , Lo $\Delta\phi$]	7000, 5000
0	[Lo Δr , Lo $\Delta\phi$], [Mid Δr , Lo $\Delta\phi$], [Mid Δr , Mid $\Delta\phi$], [Hi Δr , Hi $\Delta\phi$],	7000, 5000, 4500, 4200

viscous times at the innermost regions, but less than one viscous time at $r \gtrsim 3a$.

We have performed 28 runs altogether, for 12 different binary mass ratios between $q = 0.01$ and $q = 1.0$ (as well as control runs for $q = 0$, i.e. a single BH). The mass ratio, resolution, and the number of binary orbits followed in each of our simulation runs are summarized in Table 1.

2.3 Initial Conditions

In our initial conditions, we insert a central cavity around a binary, and also include a density pile-up just outside the cavity wall. Such a surface density profile is expected to develop during the inward migration of the secondary. When the secondary arrives at the radius where the local disk mass is too small to absorb the secondary’s angular momentum, its migration stalls, the inner disk drains onto the primary, and continued accretion from larger radii causes a pile-up of gas outside the secondary’s orbit (Syer & Clarke 1995; Ivanov et al. 1999; Milosavljević & Phinney 2005; Chang et al. 2010; Rafikov 2012). As emphasized recently by Kocsis et al. (2012a), the details of this process are still uncertain, as the coupled time-dependent migration, cavity formation, and pile-up, has not been modeled self-consistently, even in one-dimensional calculations. However, using self-consistent *steady-state* solutions, Kocsis et al. (2012b) showed that in many cases, the pile-up can cause overflow already at large binary separations.

For simplicity, and for ease of comparison, we adopt the same initial surface density profile as in MM08. This profile was motivated by the earlier results of Milosavljević & Phinney (2005); it has very little gas inside of $r \simeq 3a$, and peaks at $\sim 8a$,

$$\Sigma(r, t_0) = \Sigma_0 \left(\frac{r_s}{r} \right)^3 \exp \left[- \left(\frac{r_s}{r} \right)^2 \right]. \quad (5)$$

Here Σ_0 is an arbitrary constant, and $r_s = 10a$. The initial profile is shown by the [blue] dashed curves in Figure 4 below.

We also follow MM08, and incorporate pressure gradients and the quadrupole contribution of the binary’s poten-

tial into the initial azimuthal velocity,

$$\Omega^2 = \Omega_K^2 \left[1 + \frac{3}{4} \left(\frac{a}{r} \right)^2 \frac{q}{(1+q)^2} \right]^2 + \frac{1}{r\Sigma} \frac{dP}{dr} \quad (6)$$

and account for viscous drift in the initial radial velocities,

$$v_r = \frac{d}{dr} \left(r^3 \nu \Sigma \frac{d\Omega}{dr} \right) \left[r \Sigma \frac{d}{dr} (r^2 \Omega) \right]^{-1}. \quad (7)$$

We emphasize that with this profile, the disk is not initially in equilibrium, and material diffuses away from the peak of the surface density, both inward and outward, due to pressure gradients. However, after running the simulations for several thousand orbits, the system relaxes to a steady pattern of accretion, and we do not expect the initial profile to significantly influence our conclusions.

2.4 Tests of Code Implementation

To ensure that non-axisymmetric perturbations are not induced artificially in the disk, we simulate a standard Shakura-Sunyaev α -disk around a single black hole for 10^4 binary orbits. To keep axisymmetry during these runs, we re-implemented the FLASH2 routine UNBIASED-GEOMETRY (UBG)⁴ into the FLASH3 routines. UBG cleans up round-off errors in the cell boundary positions, and forces the polar grid to keep cell sizes uniform in the azimuthal direction. Without UBG, small perturbations in the azimuthal direction grow to significant size after a few hundred orbits. We have verified that after re-implementing UBG, axisymmetry is preserved for 10^4 binary orbits.

To test our implementation of viscosity in polar coordinates, we ran simulations of a viscously spreading ring of matter in a Keplerian potential. When the kinematic viscosity coefficient ν is constant in space and time, there exist analytic solutions for the evolution of the surface density $\Sigma(x, T)$ and radial velocity $v_r(x, T)$ (Pringle 1981). The dimensionless radial and temporal variables $x \equiv r/r_0$ and $T \equiv 12\nu r_0^{-2}$, are scaled to the initial position of the ring r_0 , and to the viscous time at r_0 , respectively.

⁴ This routine is titled *clean_last_bits* in the FLASH2 download.




Figure 1. Tests of our viscosity implementation in FLASH3.2, for a viscously spreading ring of matter in a Keplerian potential. The solid curves represent analytic solutions (Pringle 1981) and the dashed curves show the results of numerical simulations, at four different times, as labeled. $T = 12\nu t r_0^{-2}$ is the dimensionless time, scaled to the viscous time at the initial position r_0 of the ring. The top panel shows the surface density (in arbitrary units) as a function of the dimensionless radius $x = r/r_0$. The bottom panel shows the velocity profile in units of x/T .

To test our code, we replaced the potential of the binary (equation 2) by that of a point mass, and initialized FLASH with Keplerian azimuthal velocity, surface density $\Sigma(x, T_0 = 0.32)$, radial velocity $v_r(x, T_0 = 0.32)$, a constant temperature such that $v_\phi/c_s \simeq 30$ so the disk is thin, and a large, constant kinematic coefficient of viscosity such that the viscous time is of order an orbital time at the disk inner edge $t_{\text{visc}}\Omega_K(r_{\text{min}})/2\pi \simeq 1$. The latter choice of a large viscosity coefficient amplifies the effects of viscosity, and also helps avoid a small initial transient which emanates from the inner edge of the disk after ~ 1 orbital period. If this transient perturbs the initial conditions, agreement with the analytic solutions is more difficult to gauge.

Figure 1 displays the results of the test. Our implementation is in good agreement with the analytic solutions for the entire viscous timescale shown in the figure. However, we did find a peculiarity: this good agreement is achieved by choosing a value of ν in the simulations that is a factor of ~ 3 greater than used in the analytic solution. We suspect that this factor of 3 discrepancy is caused by the different FLASH2 and FLASH3 viscosity implementations, but we were unable to track down its origin. We also note

that the similar “fudge factor” of $\sim 2 - 3$ is needed for our equal-mass binary runs to agree with the accretion rates and surface densities observed in MM08. We therefore trust the results of the above tests, and simply adopt coefficients of kinematic viscosity which are 3 times larger than the desired input value (i.e., to simulate a disk with $\alpha = 0.01$, we input $\alpha = 0.03$ into FLASH3). In section § 4.3.2, we further discuss the effects on our results of changing the viscosity coefficient.

Finally, we note that the above test does not exercise any azimuthal components of the viscous momentum flux, which are present in a non-axisymmetric disk.

3 PHYSICAL REGIME

3.1 Black Hole Binary Parameters

The simulations can be scaled, in principle, to any black hole mass and orbital separation. In this section, we discuss the physical scales for which our simulations could be relevant (i.e. physically viable and observationally interesting).

In general, we are interested in primary MBHs with a mass in the range of $10^5 M_\odot \lesssim M_p \lesssim 10^9 M_\odot$. It is not clear whether smaller BHs exist in galactic nuclei, and, in any case, the radiation from such a low-mass BHs would likely be too faint to detect. Likewise, much more massive BHs are known to be rare. For the mass ratio q , the interesting range is $10^{-2} \lesssim q \lesssim 1$. As we show below, for the set-up we study (i.e. with a cavity inserted by hand into the disk), the accretion pattern converges as we decrease the mass ratio to $q = 0.01$; we expect the results for $q < 0.01$ would remain very similar. In practice, a physical lower limit of $q \sim 0.01$ may also arise from the fact that bound binary BHs can be created only in relatively major mergers. In a minor merger, the smaller satellite galaxy may not experience the torques needed to bring its gaseous nucleus, with the low-mass BH, close to the center of the larger galaxy, for the BH-BH merger to take place (Hopkins et al. 2006). Coupled with the well-established correlations between the mass of a MBH and its host galaxy, this suggests that the q -distribution may not extend to values significantly below $q \sim 0.01$.⁵

Following a merger of two MBH-harboring galaxies, the MBHs sink to the bottom of the new galactic potential via dynamical friction in approximately a galactic dynamical timescale (Begelman et al. 1980). In addition to stellar interactions (e.g. Preto et al. 2011), many studies have shown that gas in the vicinity of the binary could aid in hardening the binary down to $\ll$ pc separations (e.g. Escala et al. 2005; Dotti et al. 2007; Mayer et al. 2007; Lodato et al. 2009; Cuadra et al. 2009; Nixon et al. 2011; Chapon et al. 2011). We have assumed that such gas in the vicinity of the binary cools efficiently and forms a rotationally supported thin disk. For a given M_p and M_s , we are still free to choose a physical distance for the orbital radius a , which could correspond to a snapshot of the binary anywhere along its orbital decay. Accretion disks become self-gravitating,

⁵ In principle, less massive BHs may grow from the accretion disks around the primary (McKernan et al. 2012); the long-term evolution of such systems would be worthy of further study.

Table 2. *Orbital and residence times for binaries with total mass $M = 10^5 M_\odot$, $M = 10^7 M_\odot$, and $M = 10^9 M_\odot$, at different orbital separations a (shown in units of Schwarzschild radius r_S) and mass ratios $q = 1$ and $q = 0.01$. The residence time is defined as $t_{\text{res}} = -a(da/dt)^{-1}$, and is adopted from Haiman et al. (2009). The 5th column approximates the number of orbits until merger ($N_{\text{orb}} = t_{\text{res}}/t_{\text{orb}}$) at separation a . The last column indicates whether the orbital decay of the binary is driven by gravitational radiation or by interacting with the gas disk. Our typical simulation is run for 4500 orbits, during which the secondary would remain stationary to a good approximation.*

$M [M_\odot]$	Mass ratio q	$a [r_S]$	t_{orb}	$t_{\text{res}} [\text{yr}]$	N_{orb}	Migration
10^5	1	10^2	2.44 hr	8.0	2.9×10^4	GW
10^5	1	10^3	0.46 wk	2.0×10^4	2.3×10^6	disk
10^5	1	10^4	14.52 wk	2.2×10^5	7.9×10^5	disk
10^5	0.01	10^2	2.44 hr	2.0×10^2	7.2×10^5	GW
10^5	0.01	10^3	0.46 wk	6.2×10^3	7.0×10^5	disk
10^5	0.01	10^4	14.52 wk	5.8×10^4	2.1×10^5	disk
10^7	1	10^2	1.45 wk	8.4×10^2	6.7×10^5	GW
10^7	1	10^3	0.88 yr	6.3×10^5	7.1×10^5	disk
10^7	1	10^4	27.8 yr	5.7×10^6	2.0×10^5	disk
10^7	0.01	10^2	1.45 wk	5.7×10^3	2.0×10^5	disk
10^7	0.01	10^3	0.88 yr	1.9×10^5	2.1×10^5	disk
10^7	0.01	10^4	27.8 yr	1.5×10^6	5.1×10^4	disk
10^9	1	10^2	2.78 yr	8.0×10^4	2.9×10^4	GW
10^9	1	10^3	88.0 yr	1.7×10^7	1.9×10^5	disk
10^9	1	10^4	2.78×10^3 yr	3.0×10^8	1.1×10^5	disk
10^9	0.01	10^2	2.78 yr	1.0×10^5	3.6×10^4	disk
10^9	0.01	10^3	88.0 yr	9.5×10^6	1.1×10^5	disk
10^9	0.01	10^4	2.78×10^3 yr	3.0×10^8	1.1×10^5	disk

and unstable to fragmentation, beyond a radius of order $\gtrsim 10^4 (M/10^7 M_\odot)^{-1} r_S$ (where r_S is the Schwarzschild radius; see, e.g., Goodman 2003; Haiman et al. 2009). Since the binary has to fit inside a gravitationally stable disk, this puts an upper limit on the orbital separation. For a fiducial binary mass of $M = 10^7 M_\odot$, we have $a_{\text{max}} \sim 10^4 r_S$, or 10^{-2} pc.

Another practical consideration is the orbital time,

$$t_{\text{orb}} = \frac{2\pi}{\Omega} = 0.88 \left(\frac{M}{10^7 M_\odot} \right) \left(\frac{a}{10^3 r_S} \right)^{3/2} \text{ yr.} \quad (8)$$

As we will show, the accretion rate shows periodicity on a timescale of $\sim t_{\text{orb}}$. In a realistic survey, it will be feasible to look for periodic variations perhaps between $0.1 \text{ hr} \lesssim t_{\text{orb}} \lesssim$ few yr. Here the lower limit comes from the integration time required to measure the flux variations for MBHBs in the above mass range (for a survey instrument with a sensitivity similar to LSST; Haiman et al. 2009), and the upper limit comes from the duration of proposed time-domain surveys. For a $M = 10^7 M_\odot$ binary, this translates to $r_S \lesssim a \lesssim 10^3 r_S$, or $10^{-6} \text{ pc} \lesssim a \lesssim 10^{-3} \text{ pc}$.

For our simulations to be self-consistent, we also require $a \gtrsim 100 r_S$, since our Newtonian treatment ignores general relativity. Furthermore, at approximately the same binary separation, the orbital decay of the binary due to gravitational wave emission becomes more rapid than the viscous time at the edge of the cavity. As a result, the disk decouples from the binary and is ‘left behind’, rendering our initial conditions inconsistent in this regime (e.g. Milosavljević & Phinney 2005; although see Farris et al. 2012 and Noble et al. 2012 whose MHD simulations suggest that the gas can follow the binary down to smaller separations).

Finally, throughout our simulations, we fix the binary separation; we therefore require that the orbital decay should be slow enough for the binary’s orbit not to change significantly over a few thousand orbits. Assuming that the

binary is embedded in a thin disk, Haiman et al. (2009) compute residence times, $t_{\text{res}} \equiv -a(da/dt)^{-1}$, as a function of the binary separation a . Among the regimes they consider, the one most likely to be relevant for our binaries is the so-called ‘secondary-dominated’ type II migration, i.e. when the secondary’s mass exceeds the local disk mass. Note that there is a trade-off: a longer residence time is desirable since it increases the probability of finding such a system; however, longer residence times occur at larger separations and longer orbital times, which will make it more difficult to verify any periodic behavior.

The residence times and corresponding orbital times, for binaries with total mass $M = 10^5 M_\odot$, $M = 10^7 M_\odot$, and $M = 10^9 M_\odot$, and mass ratios $q = 1$ and $q = 0.01$, are summarized in Table 2. As this table shows, for a $10^7 M_\odot$ binary at separations of $a = 10^2 - 10^3 r_S$, the orbital time is between a $\sim$ week and a $\sim$ year, and the residence time is between 10^{5-6} years (except for the smallest separation of an equal-mass binary, which is already in the GW-driven regime, with a residence time of $\sim 10^3$ years). The binaries spend $> 2 \times 10^5$ orbits at these separations, justifying our constant a assumption over the $< 10^4$ orbits we simulate. At one extreme of the table, for $M = 10^5 M_\odot$, $q = 1$, and $a = 10^2 r_S$, the residence time of 8 years is too short for a probable detection. At the other extreme, for $M = 10^9 M_\odot$ and $a = 10^4 r_S$, the orbital time of 2780 years is much too long to pin down over a human lifetime! We also note that these results have been derived from simplified migration models; recent studies have found significantly longer residence times (Kocsis et al. 2012b; Rafikov 2012), making the above conclusions conservative.

Finally an important question is whether the binary + disk systems simulated here will indeed form a cavity during earlier stages of their evolution. For $q \geq 0.01$, our binaries certainly satisfy the standard gap-opening conditions in thin disks, requiring that binary torques balance viscous stresses

(e.g. Lin & Papaloizou 1986; Artymowicz & Lubow 1994). On the other hand, the formation of a central cavity, rather than an annular gap, also requires that the binary migration stalls, allowing the inner disk to drain. Such stalling creates the pile-up of gas outside the secondary's orbit, and depending on the details of viscosity and binary parameters, this pile-up can cause an overflow that fills the cavity. The *steady-state* solutions in Kocsis et al. (2012b, see their Figures 4 and 5) suggest that for $q \geq 0.01$, and for a $10^7 M_\odot$ binary, a cavity persists down to $a \sim 300 r_S$, before it is filled in; the overflow for smaller [larger] primary BHs occurs earlier [later].

In summary, we conclude that the intuition gained from our simulations can be applied to systems with a primary BH with $M_p \sim 10^7 M_\odot$ and binary separations $100 r_S \lesssim a \lesssim 10^3 r_S$. For such systems, the binary is in the Newtonian regime, its separation stays constant for $\gtrsim 10^5$ orbits, the disk is gravitationally stable, we expect a cavity to be present, and the orbital time-scale is conveniently in the possibly detectable range (of a week to a year). Going to more massive MBHs, the range of a allowed by requiring the disk to be stable is reduced. For $M = 10^9 M_\odot$, this leaves a relatively narrow interesting range around a few hundred r_S , where the disk is still stable, and the orbital time is a few years (see also Tanaka et al. 2012). For less massive MBHs, Toomre-stability allows larger separations, out to $a \sim 10^6 r_S$ for $M = 10^5 M_\odot$. The limitations then become long orbital times, and too few orbits spent at such large separations. Using the models of Haiman et al. (2009), and requiring $t_{\text{orb}} < 10$ yrs and at least $> 10^5$ orbits, for $M_p = 10^5 M_\odot$ we find the upper limit $a \lesssim 10^5 r_S$ (or 10^{-3} pc). Additionally, for such low-mass primary BHs, the cavity may close at relatively large radii, requiring $a \gtrsim 10^{3-4} r_S$ (Kocsis et al. 2012b). However as Table 2 shows there remains a sweet-spot for $10^5 M_\odot$ BH's at $a \simeq 10^4 r_S$, exhibiting ~ 10 week orbital periods and residence times of $\sim 10^5$ years.

3.2 Disk Parameters and Accretion Rate

Our primary goal is to quantify the magnitude and variability of accretion across the central cavity. To this end we compute the time-dependent accretion rate at the inner edge of the simulation domain, $r_{\text{min}} = a$

$$\dot{M}(t) = \int_0^{2\pi} \Sigma(r_{\text{min}}) v_r(r_{\text{min}}) r_{\text{min}} d\phi. \quad (9)$$

Since we are unable to track the fate of the gas at smaller radii, nor do we allow the masses of the BHs to increase, this mass is effectively lost from the simulation. In practice, the total mass that is lost is a small fraction of the total initial disk mass (at most a few % by the end of each run).

Because we neglect the self-gravity of the disk, equations (1) are independent of Σ_0 , and there is no unique way to assign a physical normalization to $\dot{M}_{\text{sim}}$ without appealing to a disk model which includes additional physics. Instead, MM08 compare the average accretion rate at the edge of the integration domain ($r_{\text{min}} = a$) to that in a disk in a Keplerian potential with the same surface density at $r \simeq 3a$,

$$\frac{\dot{M}_{\text{bin}}}{\dot{M}_{\text{free}}} \simeq \frac{\langle \dot{M}_{\text{sim}}(a) \rangle_t}{6\pi\alpha} \left(\frac{H}{r} \right)^{-2} (GMr)^{-1/2} \Sigma^{-1}(3a). \quad (10)$$

MM08 find $\dot{M}_{\text{bin}}/\dot{M}_{\text{free}} \simeq 0.2$. To make this comparison meaningful, one must assume that $\dot{M}_{\text{free}}(3a) \sim \dot{M}_{\text{free}}(a)$, i.e. that the reference, fictitious point-mass disk is in steady-state. However, for a steady state disk, specifying α and H/r (along with the dominant source of opacity) sets the physical value $\dot{M}_{\text{free}}$. For our fiducial values of $M = 10^7 M_\odot$ and $a = 10^3 r_S$ (and with $\alpha = 0.01$ and $H/r = 0.1$), we find the unphysically large value of $\dot{M}_{\text{free}} \sim 10^7 \dot{M}_{\text{Edd}}$ (where $\dot{M}_{\text{Edd}}$ is the accretion rate that would produce the Eddington luminosity, with a radiative efficiency of 10%). If we instead require $\dot{M}_{\text{free}} \sim \dot{M}_{\text{Edd}}$, then this would translate to a much thinner disk, with $H/r \simeq 4 \times 10^{-3}$. In such a thin/cold disk, resolving density waves with the same number of cells as in MM08 would require a radial grid resolution ~ 17 times higher than in our highest-resolution run, and would be impractical.

Rather than attempting to compare our binary simulations to a hypothetical steady-state point-mass disk, we chose to leave the surface density normalization Σ_0 essentially arbitrary, and instead performed explicit reference simulations with a single BH (i.e. $q = 0$), with the same initial conditions as the binary runs. This approach has the advantage of explicitly isolating the effect of turning on/off the binary. Note, however, that our point-mass runs should *not* be expected produce the steady-state accretion rate for a corresponding single-BH system. This is because our initial conditions are far from this state, and we do not run the simulations long enough (i.e. for a few viscous time at the outer edge) to allow it to settle to the correct steady-state.

4 RESULTS AND DISCUSSION

4.1 Equal-Mass Binary

We begin by describing the results of our equal-mass binary runs in some detail. Although these are very similar to those of MM08, this will serve as a useful point of comparison for our unequal-mass runs.

4.1.1 A Toy Model with Massless Particles

Before showing the results from our simulations, we consider a simple toy model, based on the orbits of non-interacting massless test particles around a binary. We populate a 2D disk with test particles, centered on the binary's center of mass, and leaving out a central cavity. We assign initial velocities equal to the Keplerian velocities around a single point mass $M = M_p + M_s$. We then follow the orbit of each test particle in the rotating binary potential, by numerically solving the restricted three-body problem for each individual particle (for 10^5 particles in practice, using equations 3.16 and 3.17 in Murray & Dermott 2000).

The results of this simple exercise are displayed in Figure 2, which shows the locus of the test particles initially, as well as after 0.25, 0.5, and 0.75 binary orbits (in a frame co-rotating with the binary). As this figure demonstrates, there is a tendency for the binary to pull streams of particles into the cavity. This, of course, is purely a gravitational effect. The toy-model can not be pushed much further in time, since after ~ 0.75 binary orbits, the trajectories of




Figure 2. The distortion of a disk of massless test particles, initially in circular orbits around the center-of-mass of an equal-mass binary, with a central cavity. The panels show snapshots of the locations of the disk particles after 0.25, 0.5, and 0.75 binary orbits, as labeled. The orbits of the particles were followed by solving the restricted three-body problem, and are shown in a frame co-rotating with the binary. The binary point masses are marked by the two [red] dots at $x = \pm 0.5a$. The figure illustrates the tendency of the binary to create streams of particles entering the central cavity, due to gravity alone.

the test-particles cross – this necessitates a hydrodynamical treatment. Nevertheless, the figure does suggest that an empty cavity can not be maintained by an equal-mass binary, even in the absence of pressure or viscosity. Furthermore, as we will see below, the simulations show accretion streams with morphologies quite similar to those in the bottom right panel of Figure 2.

4.1.2 Hydrodynamical Evolution: Reaching Steady State

We next present the results from our equal-mass binary simulations. The disk evolves through two distinct stages. As explained above, the disk is set up to be out-of-equilibrium, and we observe an initial transient state, which lasts for ~ 1000 orbits. We expect the details of this state to depend on the initial conditions. The disk then settles to a quasi-steady state, which persists for the rest of the simulation. In this quasi-steady state, the disk exhibits significant accretion, which varies periodically on two timescales, $(1/2)t_{\text{bin}}$ and $3t_{\text{bin}}$. We expect these latter features to be robust and insensitive to initial conditions.

Transient State. Initially, pressure forces and viscous stresses act to move material inside of the initial density peak at $r \sim 10a$ inward while pressure gradients move material outside of the density peak outward. Once the inner disk material reaches $r \simeq 2a$, its surface density becomes strongly perturbed and highly non-axisymmetric. Reminiscent of the evolution of the toy model in §4.1.1, two narrow

mirror-symmetric streams develop, in which the gas flow becomes nearly radial. About 10% of the material in these streams exits the integration domain at $r_{\text{min}} = a$. The rest gains angular momentum from the faster moving black holes and is flung back toward the bulk of the disk material at $r \simeq 2a$. This process maintains a central cavity within the disk, with gas streams being pulled in and pushed out on a period of $\sim 1.5 t_{\text{bin}}$. It is tempting to associate this period with the $\Omega_{\text{bin}} : \Omega = 3 : 2$ outer Lindblad resonance (LR), located at $r_2 = (3/2)^{2/3} a$ which is the strongest resonance available to an equal mass circular binary. However, as also emphasized in Shi et al. (2011), the usual interpretation of a LR requires the particles at resonance to be on nearly Keplerian orbits, whereas at r_2 , gas is flowing in nearly radial streams towards the binary. We offer a different geometrical explanation for the $1.5 t_{\text{bin}}$ variability below.

As more disk matter flows in from the initial density peak to the inner $r \approx 2a$ region, the streams become more dense. When these streams are flung back out and hit the opposing cavity wall, they generate noticeable overdensities, and deform the circular shape of the cavity to become elongated and lopsided. These overdensities then rotate at the disk’s orbital velocity. They spread out and propagate into the disk as differential rotation causes them to wind up, creating a mirror-symmetric ($m = 2$), rotating spiral pattern.

The disk symmetry lends an explanation for the $\sim 1.5 t_{\text{bin}}$ accretion rate variability noted above. Fluid at the cavity wall ($r \simeq 2a$), and thus the inner edge of the rotating spiral pattern, complete an orbit once every $\sim 3 t_{\text{bin}}$. Because of the $m = 2$ symmetry, the disk looks the same every $\sim 1.5 t_{\text{bin}}$. Also, the binary returns to the same state every $1/2 t_{\text{bin}}$, swapping secondary and primary. This means that for the $q = 1$ binary, the binary disk systems finds itself in the same state every $\sim 1.5 t_{\text{bin}}$ *i.e.* streams are generated once every $1.5 t_{\text{bin}}$ when the holes come nearest the lopsided disk cavity wall.

Quasi-Steady State. After the initial ~ 1000 orbits, the central cavity becomes noticeably lopsided, and stream generation becomes stronger on the near side of the cavity. This causes more mass to be driven into the cavity wall on the far side, pushing that side of the wall still farther away from the binary. Since the opposite, weaker stream is generated on the now-farther side of the cavity, it is further weakened as the disk becomes even more lopsided. This “feedback” process continues for a period of 200-300 orbits, after which the weaker stream, and its effect on the cavity wall structure disappears entirely. The central cavity takes on a steady lopsided shape, with a near-side where streams are pulled from the cavity wall by each passage of the holes, and a far side where non-accreted material from the streams is flung back and crashes into the cavity wall. At the azimuthal locations of these crashes, the far-side of the wall develops very strong shocks, with Mach numbers up to $\mathcal{M} \approx 15$. (However, since our disk is locally isothermal, this is likely an upper limit for the shock strength.)

We show examples of the two-dimensional surface density distributions in Figure 3. The top row of this figure shows snapshots at ~ 800 binary orbits (left) and at ~ 4000 binary orbits (right), of the inner 6% of the simulated disk (*i.e.* $\pm 6r/a$ are shown in both directions). The solid circle marks the inner boundary of the simulation domain at $r = r_{\text{min}} = a$. The larger dotted circle at $r \simeq 2.08a$ is




Figure 3. *Top row:* Surface density distributions for the equal-mass ratio ($q = 1.0$) binary during a transient, mirror-symmetric state after ~ 800 binary orbits (left) and during the quasi-steady asymmetric state after ~ 4000 binary orbits (right). The inset in the top left panel zooms in to the inner $\pm 2.5r/a$ of the disk (at a later orbital phase) in order to show the stream morphology. *Bottom two rows:* snapshots at ~ 4000 binary orbits, during the quasi-steady state phase, for mass ratios $q = 0.5, 0.1, 0.05$, and 0.01 , as labeled. Each panel shows the inner $\sim 6\%$ of the simulated disk, extending $\pm 6r/a$ in both directions. The solid circles mark the inner boundary of the simulation at $r = r_{\min} = a$. The larger dotted circle at $r \simeq 2.08a$ is the position of the $(m, l) = (2, 1)$ outer Lindblad resonance (shown only for reference). Surface densities are plotted with the same linear grayscale in each panel, with the darkest regions corresponding to a maximum density of $0.4\Sigma_0$ (i.e. the peak of the initial surface density profile). Orbital motion is in the clock-wise direction.

the position of the $(m, l) = (2, 1)$ outer Lindblad resonance (not present, but shown for reference). The left of these two panels illustrates the mirror-symmetric, transient state. The zoomed-in inset of this panel shows two weak mirror-symmetric streams (at a later orbital phase) reminiscent of the streams seen in the toy model of Figure 2. The right panel shows the disk after it has settled to its quasi-steady, lopsided state, with a single stream.

The top left panel of Figure 4 shows snapshots of the azimuthally averaged surface density profile of the equal-mass binary disk at three different times, after 100, 2000, and 4000 orbits. For comparison, the density profile is also shown for the single point-mass ($q = 0$) case, after 4000 orbits. As the figure shows, the inner circumbinary disk spreads inward with time. By 4000 orbits, the disk structure at $r/a \gtrsim 5$, where the effect of the binary is relatively small, closely follows the $q = 0$ case. However, the density profile remains sharply truncated inside $r \lesssim 2a$ (i.e. below the point-mass case, even at 4000 orbits). The peak density decreases as matter drains inward towards the holes and also outward towards the outer boundary. The figure also shows that the position of the cavity wall moves slightly outward between the $t=2000$ and $t=4000$ snap-shots. This is because as the disk settles to its lopsided quasi-steady state, the cavity size grows in the azimuthally averaged sense.

4.1.3 Torque Balance and the Size of the Central Cavity

As (the top left panel of) Figure 4 shows, the central cavity around the equal-mass binary extends to $r/a \approx 2$. The cavity edge is expected to be located at r_{ce} , defined as the radius where the negative viscous torque matches the binary torque,

$$\left[\left(\frac{dT}{dr} \right)_{\text{bin}} + \left(\frac{dT}{dr} \right)_{\text{visc}} \right]_{r_{ce}} = 0. \quad (11)$$

To gain insight into the transport of angular momentum and the clearing of the central cavity, we therefore compute the time- and azimuthally-averaged torque densities from the binary potential,

$$\left(\frac{dT}{dr} \right)_{\text{bin}} = \left\langle \frac{1}{2\pi} \int_0^{2\pi} \Sigma(r, \phi) \frac{d\Phi}{d\phi}(r, \phi) r d\phi \right\rangle_t, \quad (12)$$

and from viscous stresses,

$$\left(\frac{dT}{dr} \right)_{\text{visc}} = 2\pi \left\langle \frac{d}{dr} \left[r^3 \nu \left\langle \Sigma \frac{\partial \Omega}{\partial r} \right\rangle_\phi \right] \right\rangle_t. \quad (13)$$

The outer derivative in equation (13) is taken numerically, and all of the above values are measured directly from the simulation outputs, except for the binary potential derivative in equation (12) which is given analytically.

The top row of Figure 5 shows the binary and viscous torque densities for $q = 1$ over the inner $5r/a$ of the disk, during both the initial transient state (left panel) and the subsequent quasi-steady state (right panel). There is indeed a well-defined central region, where the binary torques exceed the viscous torques and can be expected to clear a cavity. The transition (computed via eq. 11) is located at $r_{ce} \simeq 1.85$ and $r = 2.05a$ in the transient and quasi-steady state, respectively, and is marked in both panels by a vertical dotted line. These vertical lines are also shown in Figure

4 and indeed lie very close to radii where the disk surface densities remain truncated.

The small outward drift of the average position of the cavity wall from the transient to the quasi-steady state is also visible in Figure 4. As the disk transitions to the quasi-steady state, the binary torque-density wavelength increases in r/a , while keeping approximately the same amplitude. These effects can be attributed to the increasingly elongated and lopsided shape of the inner cavity; when azimuthally averaged, this results in a larger cavity size.

4.1.4 Accretion Rates

The most interesting consequence of the lopsided cavity shape is on the accretion rate. In the top left pair of panels in Figure 6, we show the accretion rate, measured across the inner boundary of the simulation ($r_{\text{min}} = a$) during the quasi-steady state, after 4000 binary orbits. The upper panel shows the accretion rate as a function of time for ≈ 16 binary orbits; the solid horizontal [blue] line shows the time-averaged accretion rate during this time, and, for reference, the horizontal dashed [green] line shows the accretion rate over the same orbits in the $q = 0$ reference simulation. Interestingly, the average accretion rate onto the binary exceeds that in the single-BH case. The lower panel shows the corresponding Lomb-Scargle periodogram, measured over 100 binary orbits at a sampling rate of once per simulation time step (~ 1000 per orbit).

As mentioned above, once the quasi-steady state is reached, the accretion rate increases from that in the transient state by an order of magnitude. As Figure 6 shows, it also begins to exhibit strong (factor of ~ 2) variability. Although this figure samples the accretion only between 4000 and 4016 orbits, the pattern is remarkably steady, and repeats itself until the end of the simulation. The accretion is clearly periodic, and displays two prominent timescales, $(1/2)t_{\text{bin}}$ and $\sim 3t_{\text{bin}}$. The stronger variability at one half the orbital-time is due to the passage of each black hole by the near side of the lopsided disk and the corresponding stripping of gas streams from the cavity wall. These streams are then driven into the opposite side of the cavity (as seen in the top right panel in Figure 3; approximately 135 degrees from the generation point). The second, longer timescale corresponds to the orbital period at the cavity wall. As mentioned above, when the non-accreted material from the streams hits the far-side of the cavity, it creates an over-density, which then orbits at the disk's orbital frequency there of $2\pi(2.05a)^{3/2}(GM)^{-1/2} \approx 2.95t_{\text{bin}}$. Similar over-dense lumps have also been found and described in the 3D MHD simulations of Shi et al. (2011); more recently, Roedig et al. (2012) have also mentioned the contributions of such lumps to the fluctuations in the accretion rate.

4.2 Unequal-Mass Binaries

We next turn to the main new results of this paper, and examine the disk behaviour as a function of the mass ratio. We start with a qualitative description of how the accretion pattern changes as we decrease q , and then present a more quantitative interpretation of our results in § 4.3 below.




Figure 4. Snapshots of the azimuthally averaged disk surface density at different times (shown by different curves in each panel, from 100 to 4000 orbits, as labeled), and for different mass ratios (shown in different panels, from $q = 1.0$ to 0.01 , as labeled). In each panel, the solid [black] curve shows, for reference, the density profile in the point-mass ($q = 0$) case after 4000 orbits. The vertical dotted lines mark the radius where binary and viscous torques balance (from Figure 5); these lie close to the observed cavity edges. The vertical solid lines mark the inner edge of the integration domain ($r = r_{\min} = a$). In each case, the inner circumbinary disk spreads inward with time, but the density profile remains sharply truncated, with a low-density central cavity inside $r \lesssim 2a$.

4.2.1 Three-Timescale Regime: $0.1 < q < 1$.

The behaviour of systems with mass ratio in the range $0.1 \lesssim q < 1$ are illustrated in Figures 3, 4 and 6. These show snapshots of the 2D surface density in the quasi-steady state, the evolution of the azimuthally averaged density profile, and the time-dependent accretion rates, respectively. Additionally, in Table 3, and in the corresponding Figure 7, we show the time-averaged accretion rate as a function of q . In each case, the accretion rate is averaged over 16 orbits in the quasi-steady stage, and is quoted in units of the corresponding rate for a single-BH ($q = 0$) disk.

These figures and table illustrate several trends as q is lowered from $q = 1 \rightarrow 0.75 \rightarrow 0.5 \rightarrow 0.25 \rightarrow 0.2 \rightarrow 0.1$:

(i) The cavity becomes more compact, but less lopsided, as one naively expects when the binary torques are reduced. These effects are clearly visible in the middle row of Figure 3,

and also in the corresponding azimuthally averaged density profiles in Figure 4: the profiles look remarkably similar for $q = 0.5$ and 0.1 , except the cavity for $q = 0.1$ is smaller, and its wall is visibly sharper (as a result of azimuthally averaging over a less lopsided 2D distribution).

(ii) The secondary comes within closer proximity of the cavity wall as q is reduced. Again, this is seen in the middle row of Figure 3, and occurs both because the cavity is smaller, and because the secondary is farther from the center-of-mass of the binary.

(iii) The dense lump, created by the shocks due to the “regurgitated” stream-material thrown back out by the binary, is still present for $q = 0.5$ (see the corresponding panel in Fig. 3, between 12 and 3 o’clock at $r \sim 2a$), but is barely discernible for $q = 0.1$. Again, this trend is unsurprising - as the torques diminish, one expects weaker shocks and smaller overdensities in any resulting lump.




Figure 5. Azimuthally- and time-averaged torque density profiles in the inner disk for the equal-mass binary (top two panels) and for unequal-mass binaries (other panels, with different mass ratios q as labeled). The top left panel corresponds to the mirror-symmetric transient stage (after ~ 800 orbits) and the top right panel to the asymmetric quasi-steady stage (after ~ 4000 orbits). Only the quasi-steady stage is shown for the $q < 1$ cases. In each panel, the dashed [black] curves show the gravitational torques from the binary, and the red [dot-dashed] curves show the *negative* viscous torques. The vertical dotted line marks the radius where the viscous and gravitational torques balance (equation 11); these are close to where the azimuthally-averaged surface density profiles are found to be truncated (see Fig. 4). Time averages are taken over 25 orbits at a sample rate of 20 per orbit.




Figure 6. – continued on next page




Figure 6. The time variable accretion rate across the inner boundary of the simulation measured at $r = r_{min} = a$ (top of each pair of panels) and the corresponding Lomb-Scargle periodogram (bottom of each pair). Panels are displayed in order of decreasing binary mass ratio, starting from $q = 1.0$ at top left (on the previous page) to $q = 0.01$ at the bottom right (on this page). The average accretion rate in each panel is denoted by the solid [blue] horizontal line. The dashed [green] horizontal line in each plot shows the average accretion rate for the point mass ($q = 0$) case for reference. For $q > 0.05$, the accretion rate is strongly modulated by the binary, with either one, two, or three distinct periods present simultaneously, depending on the value of q (see text for detailed explanations). For $q \lesssim 0.05$, the binary still reduces the mean accretion rate noticeably, but does not imprint strong periodicity; the $q = 0.01$ binary is nearly indistinguishable from a single BH.

Table 3. The mean accretion rate $\dot{M}_{bin}$, averaged over 16 orbits in the quasi-steady state, for binaries with different mass ratios. The rates are shown in units of the corresponding rate $\dot{M}_{q0}$ found in a single-BH ($q = 0$) simulation. The four rows show results for different combinations of radial and azimuthal resolutions. The second row is our fiducial resolution.

q	1.0	0.75	0.5	0.25	0.1	0.075	0.065	0.06	0.05	0.04	0.03	0.01
$\dot{M}_{bin}/\dot{M}_{q0}$ [Lo Δ r, Lo $\Delta\phi$]	1.19	1.13	1.05	0.91	0.47	0.50	0.18	0.17	0.22			0.76
$\dot{M}_{bin}/\dot{M}_{q0}$ [Mid Δ r, Lo $\Delta\phi$]	1.43	1.43	1.35	1.06	0.76	0.616			0.26	0.32	0.41	0.79
$\dot{M}_{bin}/\dot{M}_{q0}$ [Mid Δ r, Mid $\Delta\phi$]					0.90							
$\dot{M}_{bin}/\dot{M}_{q0}$ [Hi Δ r, Hi $\Delta\phi$]					1.12							

(iv) The accretion streams in the $q = 0.1$ panel in Figure 3 become noticeably weaker. However, the average accretion rate, shown in Figures 6 and 7 and in Table 3, stays at approximately $\simeq 1.4\dot{M}_{q0}$ (i.e. exceeding the single-BH reference value) over the range $0.5 \lesssim q \leq 1$. For $q \lesssim 0.5$, the average accretion rate drops more rapidly, falling by nearly a factor of two to $\simeq 0.76\dot{M}_{q0}$ by $q = 0.1$.

(v) Despite a drop in the average accretion rate, the maximum accretion rate (i.e., the amplitude of the spikes in Fig. 6) *increases* as the mass ratio is decreased to $q \approx 0.1$. For example, at $q = 0.1$, the fluctuations in the accretion rate reach a factor of ~ 4 , compared to a factor of ~ 2 in the $q = 1$ case. As a result, even though the average accretion rate is $\sim 50\%$ below the $q = 1$ case, the largest spikes

reach $\sim 25\%$ *above* the corresponding maxima for $q = 1$. This could make it easier to detect and identify such binaries, despite their weaker average accretion.

(vi) Perhaps the most interesting result is shown by the Lomb-Scargle periodograms in Figure 6. As q decreases, power is clearly traded from both the $(1/2)t_{bin}$ and the $3t_{bin}$ variability timescales into the t_{bin} timescale. This is because of the increased proximity between the secondary and the cavity wall, and a corresponding larger distance between the primary and the cavity wall noted above. As a result, as q is decreased, the secondary begins to dominate the variability, pulling stronger accretion streams off the cavity wall once per binary orbit.




Figure 7. The average mass accretion rate as a function of q , and for simulations with different spatial resolution. This is the same information as in Table 3, except here shown graphically.

Focusing on the last finding: in the $0.1 \gtrsim q > 1$ case, we find that the time-dependent accretion rate displays *three distinct and sharply defined periods*, with well-defined ratios at 0.5, 1, and $3t_{\text{bin}}$. While the last of these reflects the orbit of the dense lump at $r \sim 2a$ and could depend on details of the disk properties, the first two periods are fixed by the binary alone and are independent of the disk. *The 1:2 period ratio is therefore a robust prediction; if observed, it could serve as a smoking gun signature of a binary.*

4.2.2 Single-Orbital-Timescale Regime: $0.05 \lesssim q \lesssim 0.1$.

As q is further decreased, the overall distortions to the disk become less pronounced, and approach a nearly axisymmetric, tightly wound spiral pattern (see the $q = 0.05$ panel in Figure 3). The distance between the cavity wall and the secondary further shortens, and the accretion variability becomes dominated entirely by the streams created by the secondary’s passage once per orbit. As Figure 6 shows, for $q = 0.075$ and 0.05 , the accretion rate displays a nearly perfect sinusoidal variation, with the corresponding Lomb-Scargle periodograms showing a single spike at the orbital timescale of the binary. For $q = 0.075$, the fluctuations are still large (factor of ~ 3), but by $q = 0.05$, the fluctuations strongly diminish. Interestingly, however, we find that the average accretion rate is most strongly suppressed among all of our runs for $q = 0.05$ (by a factor of ~ 5 compared to the single-BH case).

As mentioned above, the disappearance of the $(1/2)t_{\text{bin}}$ variability timescale is easy to understand qualitatively: once the primary BH remains very close to the center-of-mass, its compact orbital motion no longer impacts the disk far away. The disappearance of the $3t_{\text{bin}}$ variability timescale is also clearly attributable to the lack of any dense lump near the cavity wall for $q \lesssim 0.1$. However, the reason this lump disappears is less obvious, and warrants some discussion.

Figure 6 shows that as q decreases, the power in the $(1/2)t_{\text{bin}}$ and $3t_{\text{bin}}$ timescales decreases *in tandem*, main-

taining a roughly constant ratio. This suggests that the variability on these two timescales are causally linked. Such a causal link could arise either through the strength or the frequency of the accretion streams:

(i) As q decreases, the cavity becomes less lopsided, and the accretion rate spikes become weaker. This suggests that when these weaker accretion streams are flung back to the cavity wall, they create less overdense lumps. For $q \lesssim 0.1$, the lump may not survive shear stresses and pressure forces, and may dissolve in less than an orbital time.

(ii) As can be eyeballed from Figure 3, the stream impact zone (dense region outside the cavity) extends over an azimuth of $\Delta\phi \sim 100 - 120^\circ$. The orbital period at the cavity edge is $\sim 3t_{\text{bin}}$, implying that multiple streams can hit parts of the same lump if and only if streams are generated more than once per binary orbit.

To test whether the strength or the frequency of the streams is more important for lump generation, we repeated our simulation for an equal-mass binary, but we placed one of the BHs artificially at what would be the real binary’s center of mass. The second hole still orbits at $r = a/2$ as usual. In this setup, the cavity wall is perturbed by streams with a similar strength as in the real $q = 1$ simulation, but now only once, rather than twice per orbit. We found that this “one-armed” binary did *not* create a significant over-density at the cavity edge, nor did it excite a significant elongation of the cavity. We therefore conclude that multiple, overlapping streams are required to generate a strong lump that survives for an orbital time. This explains the roughly constant ratio of power between the $(1/2)t_{\text{bin}}$ and $3t_{\text{bin}}$ timescales: as the power at $(1/2)t_{\text{bin}}$ is reduced, the lumps become weaker, decreasing the power at $3t_{\text{bin}}$. This result also explains why the cavity becomes less lopsided as q decreases.

4.2.3 Steady-Accretion Regime: $q \lesssim 0.05$.

As we continue to decrease q from 0.05 to 0.01, we find yet another distinct regime. The overall morphology of the snapshot of a $q = 0.01$ disk in Figure 3 looks similar to the $q = 0.05$ case, except the nearly-concentric perturbations are even weaker, and the cavity still smaller. However, the similarity is quite deceptive, with the movie versions of these figures showing a striking difference.⁶ In the $q = 0.05$ case, accretion streams form and disappear periodically, but in the $q = 0.01$ case, the disk pattern becomes constant and unchanging (in the frame co-rotating with the binary). There is still a visible accretion stream, hitting the inner boundary of the simulation just ahead of the secondary’s orbit, but the stream steadily co-rotates with the binary.

As Figure 6 shows (see also Fig. 7 and Table 3), the average accretion rate has reached its minimum at $q = 0.05$. As q is further decreased, the accretion rate becomes steady, with no fluctuations, and its value *increases* back towards the $q = 0$ rate (dashed horizontal [green] line in Figure 6). Essentially, for such a light secondary, the system begins to resemble a disk with a single BH. Although a cavity is still clearly present (moving from $r \simeq 1.6a$ at $q = 0.05$ to

⁶ Movie versions of the snapshots in Figure 3 are available at <http://www.astro.columbia.edu/~dorazio/moviespage>

$r \simeq 1.4a$ at $q = 0.01$), it is being refilled, as for a single BH. In fact, after 4000 orbits, the azimuthally averaged density profiles are nearly identical for $q = 0$ and $q = 0.01$ (see Fig. 4).

Although the secondary still excites small but visible ripples in the disk (Fig. 3), by $q = 0.01$ it can no longer exert a large enough torque to pull in large streams and drive them back out to produce a lopsidedness in the circumbinary disk. The ripples are deeply in the linear regime, and they resemble the tightly-wound spiral density waves launched in protoplanetary disks (e.g. Goldreich & Tremaine 1980; Dong et al. 2011; Duffell & MacFadyen 2012), except here our background disks have a pre-imposed central cavity. In the deeply linear regime, the waves are excited by resonant interactions with the disk, and non-linear coupling to an $m = 1$ mode is no longer possible. Note the disappearance of any ripples in the azimuthally averaged surface densities for $q \lesssim 0.05$ (Fig. 4).

4.3 Summary and Discussion

In summary, we find that the behavior of the accretion rate across the circum-binary cavity as a function of q can be categorized into four distinct regimes:

(i) *Two-timescale regime*; $q = 1$. Confirming previous results, an equal-mass binary maintains a central low-density cavity of size $r \sim 2a$. However, the time-averaged accretion rate *exceeds* that for a point-mass case; there are factor of ~ 2 fluctuations around the average on two prominent time-scales, $(1/2)t_{\text{bin}}$ and $\approx 3t_{\text{bin}}$.

(ii) *Three-timescale regime*; $0.1 \lesssim q < 1$. The time-averaged accretion rate drops by a factor of ~ 2 by $q = 0.1$; however, the fluctuations continue to grow and the peak accretion rates remain high. There are three time-scales present, $(1/2)t_{\text{bin}}$, t_{bin} , and $\approx 3t_{\text{bin}}$.

(iii) *Single-orbital-timescale regime*; $0.05 \lesssim q < 0.1$. In this regime, the average accretion rate and its fluctuations both drop with decreasing q . Variability is dominated by the secondary alone, and is nearly sinusoidal on the binary period t_{bin} .

(iv) *Steady-Accretion regime*; $q \lesssim 0.05$. The accretion becomes steady, and the system overall resembles a cavity-filling single-BH disk, with small perturbations due to the secondary in the linear regime.

4.3.1 Understanding the q -Dependence of the Disk

To better understand the reason for the transitions through each regime, we note that lowering the binary mass ratio has two main consequences. First, the position of the secondary (primary) moves away from (towards) the binary's center of mass,

$$r_s(q) = a(1+q)^{-1} \quad r_p(q) = a(1+1/q)^{-1}. \quad (14)$$

Second, the size of the central cavity decreases. This is apparent visually from both the 2D surface density snapshots (Fig. 3) and the azimuthally averaged profiles (Fig. 4). The latter figure also shows the locations of the cavity edge r_{ce} expected from balancing the azimuthally averaged gravitational and viscous torques (Fig. 5), which agree well with the observed cavity sizes. In the top panel of Figure 8, we

explicitly show r_{ce} as a function of q . The points in the figure have been obtained by balancing the azimuthally averaged viscous and gravitational torques measured in the simulation (equation 11). The black line is an empirical fit to the data points at fiducial resolution, given by

$$r_{\text{ce}}(q) \simeq 3.0 \ln(q^{0.1} + 1) a. \quad (15)$$

Figure 8 shows, in particular, that for mass ratios $0.5 \lesssim q \leq 1.0$, the average cavity size stays approximately constant, while for mass ratios $q \lesssim 0.5$, the cavity shrinks more rapidly with q (cf. Artymowicz & Lubow 1994; Milosavljević & Phinney 2005). We attribute this behavior to weaker outward binary torques for smaller q . The binary torque, as a function of q and disk coordinates, can be computed from the analytic binary potential (eq. 2) and is given by

$$T_{\text{bin}}(r, \phi, q) = -r \nabla_{\phi} \Phi = -GM \frac{ar \sin(\phi - \Omega_{\text{bin}} t)}{(1+q)(1+q^{-1})} \times \left[\left(r^2 + \left(\frac{a}{1+q^{-1}} \right)^2 - \frac{2ar \cos(\phi - \Omega_{\text{bin}} t)}{1+q^{-1}} \right)^{-3/2} - \left(r^2 + \left(\frac{a}{1+q} \right)^2 + \frac{2ar \cos(\phi - \Omega_{\text{bin}} t)}{1+q} \right)^{-3/2} \right]. \quad (16)$$

We plot $T_{\text{bin}}(r, \phi, q)$ in the middle panel of Figure 8, at three constant radii in the disk, at an azimuthal coordinate $\sim 26^\circ$ behind the position of the secondary, where the outward torque is maximal. The figure shows that the outward torque, for example at $r = 2a$, increases weakly from $q = 1.0$ to 0.5 , but then decreases more rapidly, below a $1/3$ rd of its peak value, as q decreases to $q = 0.05$. Thus, the cavity size as a function of q is well correlated with the strength of the outward binary torque which clears the cavity: both are approximately constant or weakly increasing as q drops from 1.0 to 0.5 , and then both decrease rapidly as q falls towards 0 .

We can next use the apparent cavity size $r_{\text{ce}}(q)$ and the secondary's location $r_s(q)$ to estimate the impact of each hole as it passes by a fluid element in the neighboring cavity wall. We first evaluate the radial velocity imparted to the gas by the passage of the secondary, invoking the impulse approximation. This assumes that the azimuthal deceleration just before and acceleration just after the passage cancel, and only a net radial velocity component is imparted onto the fluid element. This radial velocity is approximated as the product of the radial force from the BH at closest approach, and the duration of the encounter. Here we ignore the non-axisymmetric elongation of the cavity, and take the impact parameter at the closest approach to be the distance between the secondary and the average cavity wall edge $r_{\text{ce}} - r_s$. We note that r_{ce} is a good approximation for the distance from binary center of mass to the near side of the cavity in each of our runs. This can be eyeballed by comparing the values of r_{ce} in Figure 8 to the 2D density profiles in Figure 3. We assume that the corresponding encounter time is given by

$$\Delta t_s \simeq \frac{r_{\text{ce}} - r_s}{\Omega_{\text{bin}} r_s - \Omega_{\text{ce}} r_{\text{ce}}}, \quad (17)$$

where the denominator is the relative velocity of the disk material and the passing secondary. During this encounter, the primary is passing by in the opposite direction, at a




Figure 8. Cavity size and binary torques as a function of mass ratio q . The *top panel* shows the position of the cavity wall as a function of q , in runs with different radial and azimuthal resolutions, as labeled. The points mark the radii r_{ce} at which the azimuthally averaged viscous and gravitational torques balance (equation 11). The black line is an empirical fit to these data points. The *middle panel* shows the binary torque (eq. 16) evaluated at three constant radii and at an azimuth $\sim 26^\circ$ behind the secondary, where outward torque is maximal. For $q \lesssim 0.4$, the binary torques decrease steeply with lower q . The *bottom panel*, shows an estimate of the radial velocity imparted to a disk element at the cavity wall when the secondary (blue line) and the primary (black line) pass by. The estimate is based on the empirical fit in the top panel, and on the impulse approximation.

distance $r_{ce} + r_p$, with a corresponding encounter time of

$$\Delta t_p \simeq \frac{r_{ce} + r_p}{\Omega_{bin} r_p + \Omega_{ce} r_{ce}}, \quad (18)$$

The net increase in radial velocity due to the secondary passage on the near side is then

$$\delta v_s = \frac{GM(1+q^{-1})^{-1}\Delta t_s}{[r_{ce}(q) - a(1+q)^{-1}]^2} + \frac{GM(1+q)^{-1}\Delta t_p}{[r_{ce}(q) + a(1+q^{-1})^{-1}]^2}. \quad (19)$$

while the net increase in radial velocity due to the primary passage δv_p is found by swapping p 's and s 's in equations (17) and (18).

In the bottom panel of Figure 8, we show δv_s and δv_p as a function of q . First we see that as q is decreased from 1, the impact of the primary passage by the near cavity wall decreases monotonically. This is simply due to the increase in $r_{ce} - r_p$ as q decreases, which lessens not only the radial force felt at the cavity wall but also the encounter time.

Next, notice that as q is decreased from 1 to ~ 0.4 , the impact of the secondary passage increases considerably. This is because, opposite to the primary case, $r_{ce} - r_s$ decreases with q . The most important effect of this is to increase the encounter time. Even though $r_{ce} - r_s$ decreases, the secondary's mass decreases more quickly and the radial force exerted by the secondary lessens with decreasing q . However, as the secondary moves closer to the cavity edge with decreasing q , its orbital velocity increases as $r_s^2 \propto 1/(1+q)^2$. At the same time the cavity edge moves inwards and the orbital velocity of fluid elements there also increases but at a slower rate proportional to $r_{ce}^{-1/2} \propto [3\ln(q^{0.1} + 1)]^{-1/2}$. The result is a matching of the two orbital velocities at $q \simeq 0.4$ and a formally infinite encounter time at that mass ratio. In reality, however δv_s will be cut off as a stream is generated or as the fluid element moves to larger r along the non-circular cavity edge. If we cap the encounter time at $\sim 1/2t_{bin}$ (roughly a stream generation time), then the very large δv_s near $q \sim 0.4$ is truncated. With this restriction, δv_s decreases for all $q \gtrsim 0.1$ and the secondary reaches its largest perturbing power at $q \simeq 0.1$. As q decreases below this turn around mass ratio, the impact of the secondary passage decreases as the encounter time again becomes shorter and the radial force due to the secondary continues to decrease.

The above analysis elucidates our results in the three-timescale regime and the transition to the single-timescale regime. As q decreases from 1 to 0.1, less disk fluid is pulled into the cavity over the course of a whole binary orbit, causing a drop in the time-averaged accretion rate. However, due to its closer proximity to the cavity wall and a longer encounter time with fluid elements at the cavity wall, the secondary becomes more effective at generating brief spikes in accretion (while at the same time, the effect of the primary diminishes). As a result, the maximum accretion rate, during the narrow spikes that occur once per binary orbit, increases. This latter trend changes for $q \lesssim 0.2$, where the encounter time between a disk fluid element and the secondary quickly decreases and the secondary becomes too light to generate strong accretion spikes. At around the same $q \sim 0.2$, the primary's orbit becomes too compact to generate noticeable time-variable perturbations at the cavity wall.

The impulse analysis illuminates the effect that q has on the magnitude of the induced accretion streams as well




Figure 9. The binary torques (from eq. 16), evaluated at the cavity wall edge ($r_{ce}(q)$; from eq. 15). The *top panel* shows contours of the total (primary+secondary) torque as a function of q and azimuthal angle ϕ . The secondary is located at $\phi = \pi$. Outward [inward] torques are shown in red [blue] shades. The *bottom panel* shows a one-dimensional slice of the torque per unit disk angular momentum at an azimuth indicated by the horizontal line in the top panel, just in front of the secondary ($\Delta\phi \simeq 26^\circ$). These inward torques are largest at a mass ratio of $q = 0.2$. This suggests that the strongest streams can be pulled into the cavity (and flung back out) near this binary mass ratio, as seen in our simulations.

as the relative importance of the secondary and primary for generating streams. However, the radial velocities are on the order of the relative velocities between the perturbing hole and the disk fluid element (hence stream generation), making the impulse approximation suspect. For this reason, we next investigate the azimuthal component of the acceleration imparted by the binary, by computing the torques from equations (15) and (16).

The top panel of Figure 9 shows contours of the total torque, summed over both binary components, evaluated at the cavity wall edge, as a function of q and azimuthal an-

gle ϕ . Red regions correspond to outward torques and blue regions corresponds to inward torques. The secondary is positioned at $\phi = \pi$ and moving in the direction of increasing ϕ . In the bottom panel of Figure 9, we show the torque per unit disk angular momentum at an azimuth just in front of the secondary ($\Delta\phi \simeq 26^\circ$, where inward torques are largest). These inward torques are largest at a mass ratio of $q \approx 0.2$. This suggests that the strongest streams can be pulled into the cavity (and flung back out) near this binary mass ratio. This is consistent with our findings that there is a diminished (relative to either $q = 1$ or $q = 0$) time-averaged accretion rate, but a maximum in the peak accretion rate, near $q \sim 0.2$.⁷ More generally, this is further evidence that the general trend in accretion magnitudes can be understood simply via the kinematic interaction of the binary potential with the disk, given the geometry and size of the cavity.

Figure 3 shows that the degree of elongation of the cavity decreases with mass ratio for $q \lesssim 0.5$. As we showed in §4.2.2, this is at least partly due to the decrease in stream overlap at impact, because of the diminishing strength of streams generated by the primary. However, the middle panel of Figure 8 shows that the outward torques, at positions just behind the secondary have a maximum near $q \sim 0.4$. Such outward torques are responsible for directing the streams outward, and the strength of their impact into the cavity wall. We therefore expect that the decrease in these outward torques below $q \lesssim 0.4$ also contributes to the diminishing lopsidedness of the cavity and the overdensity in the lumps that appear at the impact sites.

4.3.2 Viscosity Dependence and Comparison with MM08

Our main qualitative conclusions for the equal-mass binary case, including the morphology of the disk, and the accretion rate, are in good agreement with MM08. Nevertheless, we find two small discrepancies with MM08, which are related to different treatments of viscosity.

First, as can be seen by comparing MM08's Figure 1 with the top left panel of our Figure 4, the azimuthally averaged density of the MM08 disk near $r \sim 2a$ increases with time as the fluid builds up against the cavity wall. Roughly speaking, MM08 find that the peak surface density barely changes from its value imposed in the initial condition; only the location of the peak drifts inward, as the inner regions of the disk are accreted toward the binary, but then held up near $r \sim 2a$. In contrast, we find that the peak surface density diminishes over time; i.e. our disk drains too rapidly to allow a significant build-up in the surface density near $r \sim 2a$.

Second, comparing MM08's Figures 7 and 8 with the top left panel of our Figure 6 we see that although the

⁷ Hayasaki et al. (2007) find that the accretion rate onto the holes increases by a factor of ~ 2 (a factor of 3.25 for the secondary and no change for the primary) for a circular binary with mass ratio $q = 0.5$ compared to an equal mass binary. However, their simulations were run for only for 40 orbital times. Since the behaviour that leads to the observed accretion rates here does not arise until after ~ 1000 orbital times (and also since the accretion rate depends on gas dynamics within the region near $r \sim 1.8a$, where Hayasaki et al. (2007) inject mass as a boundary condition), it is difficult to make a comparison between the two studies.




Figure 10. Accretion rates at the inner boundary $r_{min} = a$ for the equal-mass binary as in the top left panels of Figure 6, except for a viscosity coefficient α reduced by a factor of two, from 0.01 to 0.005. The solid horizontal [blue] line in the top panel is the average accretion rate, and the bottom panel shows the Lomb-Scargle periodogram over a duration 3 times longer than displayed in the top panel (as in MM08). Comparing to Figure 6, we see that a stronger viscosity leads to a more regular accretion pattern.

magnitudes of the time-averaged accretion rates are similar, the detailed variability and the resulting Lomb-Scargle periodograms are not identical.

To investigate the cause of these discrepancies, we reran our $q = 1$ simulation with the value of the viscosity parameter α reduced by a factor of two, from 0.01 to 0.005. The accretion rate found in this case is shown in Figure 10. Comparing with the top left panels of Figure 6, we see that a weaker viscosity leads to a less regular accretion pattern, with the period at $(1/2)t_{bin}$ less dominant, and a handful of new frequencies present (with an especially strong new peak appearing at $\sim 5t_{bin}$). Comparing our lower- α results with MM08’s Figures 7 and 8, we find that the fluctuations in the accretion rate and the Lomb-Scargle periodogram match very closely those of MM08. However, in this case, the overall magnitude of the accretion is $\sim 2 - 3$ times lower than in MM08. This can be attributed to our factor of $\sim 2 - 3$ lower density near the cavity edge (which remains for $\alpha = 0.005$).

The appearance of new frequencies in the periodogram of Figure 10, compared to the top left panel of Figure 6,

can be explained by considering the Fourier-transform (FT) of each accretion time series. Imagine a train of accretion spikes similar to those in the bottom left of Figure 6. Such a time series can be constructed by convolving the shape of the individual accretion spikes with an impulse-train of Dirac-delta-functions. By the convolution theorem, the FT of this convolution is the product of the FT of the impulse train and the FT of the spike shape. The FT of an impulse train with period T is another impulse train in the frequency domain spaced by T^{-1} . The FT of the spike shape will be qualitatively similar to a *sinc* function which tapers off and rebounds as it moves away from the fundamental frequency. The width of this *sinc* function will vary inversely with the temporal width of the spike. Thus our periodogram should resemble a train of Dirac-delta-functions spaced by T^{-1} , but with amplitudes modulated by a *sinc* function. The $q = 1$, $\alpha = 0.01$ accretion time series can be considered as a product of two such impulse trains, one at $T_{1/2} = (1/2)t_{bin}$, and one at $T_3 \sim 3t_{bin}$. Likewise, the $q = 1$, $\alpha = 0.005$ case is a product of a $T_{1/2} = (1/2)t_{bin}$, and a $T_5 \sim 5t_{bin}$ train. The differences in the two periodograms can then be attributed to the different shapes of the time domain envelopes of the T_3 vs T_5 periods. The shape of the T_5 period spike (in the $\alpha = 0.005$ case) is narrower and thus the *sinc* function envelope in the frequency domain is wider, allowing us to see the ‘extra’ frequency spikes, spaced by T_5^{-1} .

In summary, we have found good overall agreement with MM08, but were unable to reproduce a perfect match to their time-dependent accretion rate for a single choice of α : we can either reproduce their periodogram (for $\alpha = 0.005$), or their time-averaged accretion rate (for $\alpha = 0.01$). We suspect that these discrepancies are caused partly by differences between FLASH2 (used in MM08) and FLASH3 (used here). In particular, the two versions implement two different routines to compute viscous fluxes. The FLASH3 viscosity routine has been updated by the method detailed in (Edgar 2006) to conserve angular momentum. More importantly, we also conclude that the details of the accretion rate variability depend on the magnitude of disk viscosity – at least for $\alpha \lesssim 0.01$. This suggests that once the two periods at t_{orb} and $(1/2)t_{orb}$ are robustly identified in an observed system, the rest of the features in the periodogram can probe disk properties, such as the viscosity. We save the full exploration of viscous effects for a future study and here continue to run all other simulations at the fiducial value of $\alpha = 0.01$.

4.3.3 Resolution Study

Up to now we have discussed the results from our fiducial set of medium-radial, low-azimuthal resolution runs ([Mid Δr , Lo $\Delta\phi$] in Table 1). Ideally, we would repeat these runs at increasingly high radial and azimuthal resolutions, until the results converge. Unfortunately, this is computationally prohibitive, and we instead chose the following approach.

(i) For the $q = 0.1$ case only, we performed two higher-resolution runs ([Mid Δr , Mid $\Delta\phi$] and [Hi Δr , Hi $\Delta\phi$] in Table 1) and one lower resolution run ([Lo Δr , Lo $\Delta\phi$] in Table 1) to look for signs of convergence. The $q = 0.1$ case is at the cusp of the three-timescale and single-timescale regimes, where

the accretion behaviour is particularly sensitive to q . We expect it to be the most sensitive to resolution effects, as well, and therefore chose this case for our resolution study.

(ii) We repeated each of our runs at the lowest resolution. Redoing the entire set of runs allows us to assess whether different q ’s are affected by resolution differently. Furthermore, at the faster speeds at lower resolution, we could afford multiple and longer runs at the cusp between the variable and non-variable regimes in order to determine the nature of this transition.

Figure 11 gives a visual impression of the surface density distribution at the four different combinations of resolutions for $q = 0.1$. While the images look similar overall, they show a clear trend: as the resolution is increased, the lumps near the cavity wall become sharper and more overdense, and the cavity becomes larger and more lopsided.⁸ This trend can be attributed to numerical dissipation when the regurgitated streams impact the cavity wall. Increasing the resolution implies weaker numerical diffusion, stronger accretion streams, more momentum carried into the disk, and an overall more efficient driving of the $m = 1$ mode.

Figure 12 shows the corresponding time-dependent accretion rates at the inner boundary of the simulation. The accretion pattern look visibly different for the lowest resolution run which exhibits a maximum accretion rate ~ 3 times lower than its higher resolution counterparts. However, encouragingly, the difference between our fiducial [Mid Δr , Lo $\Delta\phi$] and the highest-resolution [Hi Δr , Hi $\Delta\phi$] cases is modest. The differences in the Lomb-Scargle periodograms can again be explained by considering the Fourier-transform of each accretion time series as discussed above in §4.3.2.

Table 3 and the corresponding Figure 7 shows the time-averaged accretion rates as a function of q at different resolutions. In all cases, at the same fixed q , we find that increasing resolution produces a higher accretion rate. This is consistent with the interpretation above that better resolution allows stronger accretion streams. Interestingly, we find a strong correlation between the values of $\dot{M}_{\text{bin}}/\dot{M}_{q0}$ listed in Table 3 and the accretion patterns seen in Figure 12: runs at different resolutions but with similar values of $\dot{M}_{\text{bin}}/\dot{M}_{q0}$ have very similar accretion patterns (including the variability and the values of the maxima). The result of increasing [decreasing] the resolution can therefore be interpreted as a shift of the accretion behavior to lower [higher] mass ratios.

Encouragingly, the two higher azimuthal resolution runs at $q = 0.1$, also plotted in Figure 7, lie closer to each other than the two lower resolution runs. Since the resolution steps are evenly spaced, we consider this evidence that the simulations are converging monotonically with resolution.

To find whether there is a cut-off mass ratio between variable and non-variable accretion (or whether our simulations do not run long enough to capture variable behavior for small q) we have run a number of low-resolution simulations spaced by $\Delta q = 0.005$ (see Table 3). At our fiducial value of α we find the transition from periodically variable to steady accretion occurs at $q \simeq 0.06$. At the mass ratio of

$q = 0.065$, the disk becomes eccentric, and causes variable accretion after only 3000 binary orbits. The $q = 0.06$ case is run for 8000 orbits and never becomes eccentric.

4.3.4 Caveats

However instructive, the simulations presented here are still of course simplified models of a real binary disk system, and it is worth listing some major caveats.

(i) Our simulations are two dimensional – we expect that the 3D vertical structure could modify the structure of the accretion streams, including their q -dependence.

(ii) Our simple α -viscosity prescription may be inaccurate, especially in the nearly radial accretion streams. Recent MHD simulations in the $q = 1$ case find a much larger effective α than we adopted here (Shi et al. 2011; Noble et al. 2012). We have argued that much of the accretion behavior can be understood by gravitational kinematics alone, but this needs to be verified explicitly, especially in light of the α -dependence of our results.

(iii) We assumed a locally isothermal equation of state; more realistic equations of state could have an especially large impact on the strength and dissipation of shocks at the cavity wall.

(iv) Our initial conditions correspond to an unperturbed, near-Keplerian, circular disk, and circular binary orbit, with a significant pile-up of gas. In reality, accretion onto the binary could produce significant binary (as well as disk) eccentricity (*e.g.* Cuadra et al. 2009; Lodato et al. 2009; Roedig et al. 2011). In this case, the variability we found would most likely have a more complex structure (*e.g.* Hayasaki et al. 2007) due to the plethora of resonances available in an eccentric binary potential (*e.g.* Artymowicz & Lubow 1994).

(v) Although we begin with an “empty” cavity, this cavity may overflow already at a large radius (Kocsis et al. 2012b). Future studies should construct a self-consistent initial density profile, by evolving the binary’s orbit from large radius through gap clearing, and thus determining whether a true pile-up occurs.

(vi) We have ignored the radiation from the gas accreted onto the BHs. Given that we find high accretion rates – comparable to those for a single BH – the secondary BH can be fed at super-Eddington rates, and the flow dynamics can then be strongly affected by the radiation.

(vii) We have not allowed accretion onto the BHs, and have excised the inner region $r < a$ from the simulation domain. This could have an impact on the dynamics of the streams that are flung back towards the cavity wall, and therefore on the formation of the dense lump, the lopsidedness of the cavity, and the variable accretion patterns.

These caveats should all be pursued in future work, to assess the robustness of our results. We expect that our main conclusions, namely that the accretion rate is strongly modulated by the binary, and that the power-spectrum of the accretion shows distinct periods, corresponding to the orbital periods of the binary and the gas near the location of the cavity wall, will be robust to all of the above caveats. However, the numerical values, such as the mean accretion rate, and the critical value of q for the transition between variable and steady accretion, will likely be affected.

⁸ Figure 8 shows that the *average* position of the cavity wall, found from the azimuthally averaged torques in equation (11), is largely unaffected over the range of resolutions studied here.




Figure 11. Two-dimensional surface density distributions at 4000 orbits, as in Figure 3, except for the single mass ratio $q = 0.1$, and for four different combinations of high/low radial and azimuthal resolutions, as labeled. Increasing the spatial resolution decreases numerical diffusion, leads to sharper features, and allows stronger accretion streams. The stronger streams lead to more overdense lumps where the regurgitated streams hit the cavity wall. As a result, the cavity becomes larger and more lopsided as the resolution is increased.

5 CONCLUSIONS

We have investigated the response of an accretion disk to an enclosed binary via two-dimensional, Newtonian, hydrodynamical simulations. As previous work has shown (Artymowicz & Lubow 1994; Hayasaki et al. 2007; MM08; Cuadra et al. 2009; Shi et al. 2011; Roedig et al. 2012), for non-extreme mass ratios, the binary carves out a cavity in the disk, but gas still penetrates the cavity in streams which possibly accrete onto the binary components. Here we have followed up on the work of MM08 by investigating the nature of this inflow across the circumbinary cavity, as a function of binary mass ratio q . We have simulated 12 different mass ratios in the range $0.01 \leq q \leq 1$. This corresponds to the expected range of q -values for massive BH binaries produced in galaxy-galaxy mergers.

We find that while the binary ‘propellers’ are effective at maintaining a low-density cavity at the center of the disk, they can not efficiently suppress accretion across the cavity. For $q \sim 1$, the accretion rate can even modestly exceed the rate for a corresponding single BH. The largest suppression, by a factor of ~ 5 compared to a single-BH disk, occurs for $q \approx 0.05$. As long as the circumbinary disk is fueled at a near-Eddington rate from large radius, these binaries could therefore have quasar-like luminosities. This should facilitate finding counterparts to GW events (Kocsis et al. 2006), and should also allow their detection in electromagnetic surveys (Haiman et al. 2009).

We have found that the accretion is not only strong, but can be strongly variable, with a characteristic q -dependent frequency pattern. While the accretion for $q < 0.05$ is steady,




Figure 12. The time-dependent accretion rates as in Figure 6, except for the single mass ratio $q = 0.1$ and the four different resolutions also shown in Figure 11. The $q = 0.1$ binary is at the cusp of the transition from the three-timescale to the single-timescale regime, and is particularly sensitive to resolution, as seen especially in the maxima of the accretion spikes.

for $q \gtrsim 0.05$ there is a strong modulation by the binary, and a clear dependence on q of both the variability pattern, and the magnitude of the time-averaged accretion rate. For an equal mass binary, the accretion rate is modulated at twice the orbital frequency and $\sim 1/3$ the orbital frequency. As the mass ratio is lowered, the power in the $1/2t_{\text{bin}}$ and $3t_{\text{bin}}$ variability timescales is reduced, and traded for a third variability timescale at t_{bin} . In the range $0.05 \lesssim q \lesssim 0.1$, only the single t_{bin} timescale is present. Interestingly, even when the time-averaged accretion rate is suppressed (compared to $q = 1$), we find that the fluctuations increase, and the accretion rate during the brief maxima is enhanced. We were able to interpret most of the above results based on simplified models of the binary torques and kinematics.

The strong and highly variable accretion, with characteristic frequencies, should aid in identifying massive BH binaries in galactic nuclei. The presence of two frequencies, in the ratio 1:2, is an especially robust prediction that is *independent of disk properties*, and could serve as a ‘smoking gun’ evidence for the presence of a binary. Our results suggest that the ratio of the power at these two frequencies could probe the mass ratio q , while other features of the periodogram could probe properties of the disk, such as its viscosity.

The variability time-scales are on the order of the orbital period, and we have argued that the most promising candidates in a blind electromagnetic search would be $10^6\text{--}10^7 M_\odot$ binaries, with orbital periods of days to weeks. The time-variable accretion to the central regions could produce corresponding variability in broad-band luminosities, allowing such a search in a large time-domain survey, such as LSST, without spectroscopy. Additionally, the emission lines could exhibit periodic shifts in both amplitude and frequency; kinematic effects from the binary’s orbit could be distinguished from those due to the fluctuating accretion rate, whenever the latter contains multiples of the binary period.

Parts of the accretion streams generated periodically fuel the BHs, but part of the stream material is flung back and hits the accretion disk farther out. The shocks produced at these impact sites are prominent for $q \gtrsim 0.1$, and can provide additional observable signatures. In particular, radiation from these shocks should be temporally correlated with the luminosity modulations arising near the secondary and/or primary BH, with a delay time on the order of a binary orbit.

GW observatories, such as eLISA, will be able to constrain the mass ratios of in-spiraling MBHB’s at the centers of galactic nuclei ab-initio, providing a template for the expected variability pattern. This should be helpful in identifying the unique EM counterpart among the many candidates (with luminosity variations) in the eLISA error box, as the source with a matching period.

In summary, our results imply that massive BH binaries can be both bright and exhibit strongly luminosity variations, at the factor of several level. This raises the hopes that they can be identified in a future, suitably designed electromagnetic survey, based on their periodic variability. Although encouraging, these conclusions are drawn from simplified 2D hydrodynamical models of a real binary disk system, and should be confirmed in future work.

ACKNOWLEDGMENTS

We thank Bence Kocsis for useful discussions. We also thank Paul Duffell for useful discussions and for verifying some of our results in independent runs with the code DISCO. We acknowledge support from NASA grant NNX11AE05G (to ZH and AM). DJD acknowledges support by a National Science Foundation Graduate Research Fellowship under Grant No. DGE1144155.

REFERENCES

- Amaro-Seoane P., et. al 2012, ArXiv e-prints
- Artymowicz P., Lubow S. H., 1994, *Astrophysical Journal*, 421, 651
- Barnes J. E., Hernquist L., 1992, *ARA&A*, 30, 705
- Begelman M. C., Blandford R. D., Rees M. J., 1980, *Nature*, 287, 307
- Chang P., Strubbe L. E., Menou K., Quataert E., 2010, *MNRAS*, 407, 2007
- Chapon D., Mayer L., Teyssier R., 2011, *MNRAS*, submitted, e-print ArXiv:1110.6086
- Cuadra J., Armitage P. J., Alexander R. D., Begelman M. C., 2009, *Monthly Notices of the Royal Astronomical Society*, 393, 1423
- Dong R., Rafikov R. R., Stone J. M., 2011, *ApJ*, 741, 57
- Dong R., Rafikov R. R., Stone J. M., Petrovich C., 2011, *ApJ*, 741, 56
- Dotti M., Colpi M., Haardt F., Mayer L., 2007, *Monthly Notices of the Royal Astronomical Society*, 379, 956
- Duffell P. C., MacFadyen A. I., 2012, *ApJ*, in press, e-print arxiv:1202.5608
- Edgar R. G., 2006, ArXiv Astrophysics e-prints
- Eracleous M., Boroson T. A., Halpern J. P., Liu J., 2011, ArXiv e-prints
- Escala A., Larson R. B., Coppi P. S., Mardones D., 2005, *The Astrophysical Journal*, 630, 152
- Farris B. D., Gold R., Paschalidis V., Etienne Z. B., Shapiro S. L., 2012, ArXiv e-prints
- Farris B. D., Liu Y. T., Shapiro S. L., 2011, *PRD*, 84, 024024
- Ferrarese L., Ford H., 2005, *SSR*, 116, 523
- Fryxell B., Olson K., Ricker P., Timmes F. X., Zingale M., Lamb D. Q., MacNeice P., Rosner R., Truran J. W., Tufo H., 2000, *The Astrophysical Journal Supplement Series*, 131, 273
- Gergely L. Á., Biermann P. L., 2012, ArXiv e-prints
- Goldreich P., Tremaine S., 1980, *Astrophysical Journal*, 241, 425
- Goodman J., 2003, *Monthly Notice of the Royal Astronomical Society*, 339, 937
- Haehnelt M. G., Kauffmann G., 2002, *MNRAS*, 336, L61
- Haiman Z., Kocsis B., Menou K., 2009, *The Astrophysical Journal*, 700, 1952
- Haiman Z., Kocsis B., Menou K., Lippai Z., Frei Z., 2009, *Classical and Quantum Gravity*, 26, 094032
- Hayasaki K., Mineshige S., Sudou H., 2007, *Publications of the Astronomical Society of Japan*, 59, 427
- Hopkins P. F., Hernquist L., Cox T. J., Di Matteo T., Robertson B., Springel V., 2006, *ApJS*, 163, 1
- Ivanov P. B., Papaloizou J. C. B., Polnarev A. G., 1999,

- Monthly Notices of the Royal Astronomical Society, 307, 79
- Kocsis B., Frei Z., Haiman Z., Menou K., 2006, *ApJ*, 637, 27
- Kocsis B., Haiman Z., Loeb A., 2012a, ArXiv e-prints
- Kocsis B., Haiman Z., Loeb A., 2012b, ArXiv e-prints
- Kocsis B., Haiman Z., Menou K., 2008, *ApJ*, 684, 870
- Komossa S., 2006, *Memorie della Società Astronomica Italiana*, 77, 733
- Kormendy J., Richstone D., 1995, *ARA&A*, 33, 581
- Lin D. N. C., Papaloizou J., 1986, *Astrophysical Journal*, 307, 395
- Lippai Z., Frei Z., Haiman Z., 2009, *ApJ*, 701, 360
- Lodato G., Nayakshin S., King A. R., Pringle J. E., 2009, *MNRAS*, 398, 1392
- Lommen A. N., 2012, *Journal of Physics Conference Series*, 363, 012029
- MacFadyen A. I., Milosavljević M., 2008, *The Astrophysical Journal*, 672, 83
- Mayer L., Kazantzidis S., Madau P., Colpi M., Quinn T., Wadsley J., 2007, *Science*, 316, 1874
- McKernan B., Ford K. E. S., Lyra W., Perets H. B., 2012, *ApJ*, in press, e-print arXiv:1206.2309
- Milosavljević M., Phinney E. S., 2005, *The Astrophysical Journal*, 622, L93
- Murray C. D., Dermott S. F., 2000, *Solar System Dynamics*. Cambridge University Press; Cambridge, UK
- Nixon C. J., Cossins P. J., King A. R., Pringle J. E., 2011, *Monthly Notices of the Royal Astronomical Society*, 412, 1591
- Noble S. C., Mundim B. C., Nakano H., Krolik J. H., Campanelli M., Zlochower Y., Yunes N., 2012, *ApJ*, submitted, e-print arXiv:1204.1073
- Preto M., Berentzen I., Berczik P., Spurzem R., 2011, *The Astrophysical Journal Letters*, 732, L26
- Pringle J. E., 1981, In: *Annual review of astronomy and astrophysics*. Volume 19. (A82-11551 02-90) Palo Alto, 19, 137
- Rafikov R. R., 2012, *ApJ*, submitted, e-print arXiv:1205.5017
- Robertson B., Cox T. J., Hernquist L., Franx M., Hopkins P. F., Martini P., Springel V., 2006, *ApJ*, 641, 21
- Roedig C., Dotti M., Sesana A., Cuadra J., Colpi M., 2011, *MNRAS*, 415, 3033
- Roedig C., Sesana A., Dotti M., Cuadra J., Amaro-Seoane P., Haardt F., 2012, ArXiv e-prints
- Sesana A., Haardt F., Madau P., Volonteri M., 2005, *ApJ*, 623, 23
- Sesana A., Roedig C., Reynolds M. T., Dotti M., 2012, *MNRAS*, 420, 860
- Shakura N. I., Sunyaev R. A., 1973, *Astron. Astrophys.*, 24, 337
- Shen Y., Loeb A., 2010, *ApJ*, 725, 249
- Shi J.-M., Krolik J. H., Lubow S. H., Hawley J. F., 2011, ArXiv e-prints
- Springel V., Di Matteo T., Hernquist L., 2005, *ApJL*, 620, L79
- Syer D., Clarke C. J., 1995, *Monthly Notices of the Royal Astronomical Society*, 277, 758
- Tanaka T., Menou K., Haiman Z., 2012, *MNRAS*, 420, 705
- Volonteri M., Haardt F., Madau P., 2003, *ApJ*, 582, 559